

"The world's first intelligent pain assessment tool"

Adeel Arshad (MPharm, PIP)
Omnicare Pharmacy/PainChek UK



Problem Of Managing Pain In Dementia

Pain is common

- Up to 80% of people in aged care experience chronic pain¹
- More than 70% of people in aged care have dementia²

Over 50% of people living with dementia have undetected and untreated pain¹

- Pain is difficult to assess
- Tools to assess pain are often not used or use subjective rating scales
- Difficult to use existing methods in practice
- Adverse impact on quality of care, operational efficiency and compliance





¹ Aged Care Awareness

² CPEC London School Of Economics Nov 19

³ Carter, D (2015) Dementia and Homecare: Driving Quality and Innovation by the UK Homecare Association

Behavioural Symptoms



>90% of people with dementia affected by BPSD, often caused by pain.

Greater the pain = Greater BPSD & severity

Pain group 3.8 times more likely to experience BPSD

25.3% more neuropsychiatric behaviours

33.6% higher severity of behaviours


31.4% more distress to caregivers








The PainChek Solution






Pain Domains

PainChek assesses pain across 6 key areas. These are the:

- -Face
- -Behaviour
- -Voice
- -Activity
- -Movement
- -Body





Learnings from implementation







- Improved diagnosis
- Improve BPSD
- Reduce use of antipsychotics
- Improve use of analgesics
- Preventing falls
- Reduce safeguarding incidents
- Complete pain profile

- Provide training
- Solve documentation issues
- Maintain interaction with residents
- Improve communication
- Repurposing staff
- Empower staff/staff confidence
- Influencing positive change
- Saves time







Adeel Arshad

Clinical Services and Training Manager

Mobile: 07741 193742

Email: adeel@omnicarepharmacy.co.uk

Any questions?

