How to prepare Flucloxacillin 1g

Information

1.	Diluent for reconstitution:	Water for Injection
2.	Volumes for reconstitution:	1g vial = 20ml
3.	Administration:	Inject slowly over 3 - 4 minutes
4.	Storage requirements:	Protect unopened vials from light and heat and store below 25^{o}C

Do not mix with tobramycin.

Equipment

20ml Water for Injection
Flucloxacillin vial 1g
10ml syringe(s)
Green needle(s)
Venting needle (optional)
Sterile alcohol wipes

Procedure

1. Check drug, dose and expiry date
2. Wash hands thoroughly
3. Ensure preparation area is clean
4. Swab top of vial with a sterile alcohol wipe and allow to dry for one minute
5. Take ampoule(s) of water for injection and open
6. Using 10ml syringe(s) and green needle(s) withdraw 20ml of water for injections
7. Inject water for injections into vial of flucloxacillin
8. Shake vial to dissolve powder and until solution turns clear. Make sure no particles are present (If any particles are present, discard the vial)
9. Place venting needle into vial (optional)
10. Using 10ml syringe(s) and green needle(s) withdraw contents of vial(s)
11. Tap syringe gently to allow any air bubbles to travel to the top of the syringe(s)
12. Remove any air bubbles in syringe(s)
13. Label syringe(s)

Douglas McCabe, Pharmacist
August 2010