



Influenza – Guidance for Care Homes

Health Protection Scotland

Version 1.0: October 2018





Contents

Intr	oduction	3
1.	Roles and Responsibilities:	4
2.	Key information for Care Home staff	5
3.	Checklist to prepare for influenza season	7
4.	Influenza Outbreak Checklist	8
5.	Influenza Outbreak Daily Actions Checklist	.10
6.	Influenza Outbreak Data Record	.11
7.	Influenza - Information for residents and visitors	.13
8.	Posters for use during outbreaks	.15





Introduction

The Influenza Guidance for Care Homes has been developed to provide key information to assist with prevention and control of cases of influenza in care homes. The checklists included in this pack should be used in addition to the Standard Infection Control Precautions (SICPs) and Transmission Based Precautions (TBPs) as per the <u>National Infection Prevention and Control Manual (NIPCM)</u>. This pack comprises:

- Roles and Responsibilities to guide decision making
- Information for Care Home staff
- A prevention checklist to be completed prior to the season starting
- A checklist for everything that needs to be done on the first day an outbreak is identified
- A daily checklist for every day until an outbreak is over
- An outbreak data record sheet for tracking cases and symptom onsets
- Information for visitors and residents
- Posters for displaying during the season and during outbreaks

Influenza should be suspected in any person who develops fever/high temperature in addition to acute respiratory symptoms

If 2 or more people have these symptoms within 2 days of each other, IT COULD BE AN OUTBREAK

IF YOU CONSIDER AN OUTBREAK OF INFLUENZA IS LIKELY:

- Contact your local health protection team (HPT) immediately
- Start the Influenza Outbreak Checklist





1. Roles and Responsibilities:

Person in charge of Care Home
Ensure ongoing compliance with Standard Infection Control Precautions (SICPs) and Transmission Based Precautions (TBPs).
Report all residents with acute respiratory infections (i.e. suspected influenza) to the GP.
Report to your HPT if you and the GP are clinically suspicious of 2 or more cases of acute respiratory infection (i.e. influenza) with onset within 2 days of each other.
\Box Work with the HPT in completing the daily assessments and keeping the HPT updated.
\Box Ensure there are sufficient staff available to deal with the residents' needs.
GP
Provide clinical assessment to determine outbreak status.
Provide advice to person in charge of Care Home to inform HPT.
□ Provide specific advice regarding treatment options for residents and staff.
Health Protection Team (HPT)
Advise on the control measures required and specimens to be collected.
□ Offer specific advice regarding treatment and vaccination protocols for residents and staff.

Advise on discontinuation of control measures and actions prior to re-opening of the Care Home if closed at any time





2. Key information for Care Home staff

What is Influenza?

Influenza or 'Flu' is an acute viral infection affecting the respiratory tract.

There are three main types of influenza virus (A, B & C) that affect humans but Influenza A is the most common and usually the cause of flu outbreaks in winter.

What are the symptoms?

Influenza should be suspected if the following symptoms are present:

A FEVER/HIGH TEMPERATURE PLUS ANY OF THE FOLLOWING:

- Loss of physical functioning/mental decline
- Sore throat
- Shortness of breath (more than usual)
- Productive cough (with or without sputum)
- Aching muscles

- Other chest symptoms/ chest pain
- Hoarseness/wheezing
- Runny nose or congestion
- Increased lethargy and/or general malaise
- Loss of appetite/reduced food intake

• Headache

Residents should be closely monitored for relevant signs and symptoms, particularly during the winter months when flu is most common.

If you suspect a resident has flu, IMMEDIATELY consult the resident's GP and the person in charge

How long do symptoms last?

For most people, symptoms usually resolve within 3-7 days.

Care home residents are extremely vulnerable because they are often elderly and frail, have underlying health conditions and are living in a closed environment where the virus can spread easily. Serious health complications, including secondary chest infections and pneumonia, can develop as a result of the flu and in these situations it can be life-threatening.

How is it treated?

Healthy people usually recover with rest, plenty of fluids, and over-the-counter paracetamol or ibuprofen. People at greater risk, including care home residents, may be given prophylactic antiviral medication, regardless of whether they have had their seasonal flu vaccination. Some people may require supportive care in hospital.





How is it spread?

Influenza virus is spread by **respiratory droplets**. Infected people cough or sneeze and the virus is sprayed into the air, landing on nearby surfaces and objects or into their hands. It is passed to other people when they are in close contact i.e. they are touching hands or are sneezed on, or when they touch surfaces or objects that are contaminated with droplets. **People can be infectious 1 day before their symptoms begin until 5-7 days after their symptoms have gone however the average infectious period is 2 days**.

What is an influenza outbreak?

An outbreak is when 2 or more people develop flu symptoms with onset within 2 days of each other within the same healthcare setting.

Confirmed Case	Any resident who has/has had an influenza like illness and has tested positive for influenza
Possible case	Any patient with an influenza like illness not yet laboratory confirmed
Asymptomatic case	Any patient who is asymptomatic but has tested positive for influenza

How can you reduce the risk of an influenza outbreak happening?

- Getting the flu vaccine is the single best way to protect against catching and spreading the virus it is modified each year to be as effective as possible.
- Refusing to be vaccinated puts yourself, your family, and the people you care for at unnecessary risk.
- Care home employers are responsible for ensuring vaccination arrangements are in place for their care home staff who have direct contact with residents.

What should you do if you suspect an outbreak of influenza in your care home?

- ✓ Inform the person in charge and contact the resident's GP for an assessment.
- Symptomatic residents should be cared for in their own rooms with the door closed until recovered with minimal contact with other residents e.g. avoid communal areas, have meals in their room etc.
- Use the appropriate personal protective equipment (disposable gloves, aprons, masks) when carrying out direct care.
- Wash your hands frequently with liquid soap and water as per the WHO 5 Moments; alcohol based hand rub (ABHR) can be used between washes.
- Practice good respiratory/cough hygiene.
- Encourage and assist residents and visitors to practice good respiratory/cough hygiene and good hand hygiene.
- Decontaminate frequently touched surfaces (door handles, light switches, call bells etc.) at least once daily and as often as you can during your shift, using a cleaning agent containing 1000 parts per million available chlorine.
- ✓ Make sure environmental and equipment cleaning schedules are in place and followed.
- Use the Influenza Outbreak Checklist and the Influenza Outbreak Daily Actions Checklist to organise infection prevention and control.





3. Checklist to prepare for influenza season

Act	Actions to prepare for influenza season									
1	Staff have been provided with information about the seasonal influenza	vaccination.								
2	All staff involved in patient care have received this year's seasonal flu vaccine.									
3	Influenza vaccinations for residents have been organised with the Care Home GP(s).									
4	The flu vaccination status of staff is recorded and is kept up to date. (which will increase staff safety and inform care delivery during an outbreak)									
5	This guidance document has been read by all staff members.									
6	Staff education has been reinforced for respiratory and hand hygiene.									
7	Liquid soap and disposable paper hand towels are available in all public and communal toilets and in residents' en-suites.									
8	Sufficient quantities of alcohol based hand rub (ABHR) and disposable tissues are available for public and communal areas, and resident rooms.									
9	Waste disposal systems including foot operated bins are in place throughout the facility.									
10	Personal Protective Equipment (PPE) is available (disposable gloves, aprons and masks) in sufficient quantities and stored in a clean/dry area until required for use.									
11	A segregation system is in place for infectious laundry and water-soluble bags are available.	e laundry								
12	Sufficient quantities of cleaning materials are available including a clean that contains 1000 parts per million available chlorine.	ing agent								
13	Appropriate managures are queilable for residents with symptoms for a minimum of									
14	HPS Influenza Guidance for Care Homes poster are displayed in highly	visible areas.								
15	'Catch it, Bin it, Kill it' posters are displayed in highly visible areas.									
Υοι	Ir local Health Protection Team contact details:									
	Completed by: Date:									
Des	signation:	Designation:								





4. Influenza Outbreak Checklist

Checklist to be completed as soon as an outbreak is suspected.

Date the outbreak was identified: RESIDENT PLACEMENT	Please tick $$:
Ask symptomatic residents to stay in their rooms; explain the need to keep the door of	
If a resident cannot stay in their room, carry out a risk assessment and consider other options	r care
Inform relatives and visitors of the isolation requirements and additional infection continues	trol
ADMISSIONS, TRANSFERS, DISCHARGES	
Following HPT advice, delay any planned new admissions to the Care Home	
Consider temporary suspension of visiting with the HPT if considered beneficial to ga control	ining
Following HPT advice, suspend inter-care transfers/medical appointments unless clin essential	ically
Following HPT advice, notify all receiving units (ambulances/hospitals) of the outbrea resident has to be transferred	k if a
STAFF	
Ensure that all staff on duty are asymptomatic; send all symptomatic staff home and opermit to return to work when they have been symptom-free for a minimum of 24 hou	
If possible, allocate staff to care for either symptomatic or non-symptomatic residents both	but not
If possible, allocate staff who have had their influenza vaccine to care for symptomati	с
Postpone visits from non-essential providers i.e. hairdressers, podiatrists etc.	
VISITORS	
Visitors with flu symptoms should be asked to not visit until they have been symptom- at least 48 hours	-free for
Discuss temporary suspension of visiting with the HPT if considered beneficial to gair control	ning
Offer visitors fluid resistant surgical facemasks (FRSM) if visiting a symptomatic resid the visitor has an underlying health condition and/or is at risk of more severe infection become ill	
CLINICAL MANAGEMENT	
Ensure symptomatic residents are clinically assessed by a GP (monitor continuously signs of respiratory distress (i.e. shortness of breath), dehydration, or further health deterioration)	for any
Following HPT advice, consider offering residents and staff the influenza vaccine if no already vaccinated; and/or antiviral medication to residents	ot
Following HPT advice, take samples from any resident with relevant symptoms for lal testing	ooratory
HAND HYGIENE (HH) & PERSONAL PROTECTIVE EQUIPMENT (PPI	E)
Use alcohol based hand rub (ABHR) if hands are clean, otherwise wash with liquid so water	ap and
Disposable aprons, gloves and fluid resistant surgical masks (FRSM) to be worn for e care episode	
Surgical masks should be removed and disposed of inside the patient room when at I feet (1 metre) from the resident(s)	east 3



CARE ENVIRONMENT

Remove fans and any other equipment that could increase environmental contamination

De-clutter the environment to allow effective cleaning

Remove exposed food stuffs from resident rooms and communal areas

Provide tissues and covered sputum pots for residents; dispose of tissues as healthcare waste

Provide tissues and foot-operated bins for the disposal of used tissues in communal areas

Clean all frequently touched surfaces (i.e. door handles, light switches) preferably at least twice daily using a cleaning agent that contains 1000 parts per million available chlorine (ppm av cl.)

A terminal clean of residents rooms performed once resident is no longer considered infectious and/or following resident discharge

A terminal clean of the wider facility performed once all residents and staff have been symptom-free for ≥48 hours (i.e. once the outbreak is over, with HPT agreement)

CARE EQUIPMENT

Provide resident-dedicated care equipment if possible i.e. commodes, washbowls, lifting equipment

Clean communal resident equipment between residents using a cleaning agent with 1000 ppm av cl.

COMMUNICATIONS & KNOWLEDGE

Ensure all staff members (Carers, cleaners, catering, bank staff) are aware of the situation, their responsibilities and what to do if they or a resident develop symptoms

Place appropriate signage in highly visible places at the entry to the home and in affected areas

Inform all visitors/relatives of the situation, precautions and risks- particularly for vulnerable groups

Provide relatives/visitors with Washing Clothes at Home Leaflet

Provide relatives/visitors with 'Influenza - Information for residents and visitors'

Further information on Standard Infection Control Precautions (SICPs) and Transmission Based Precautions (TBPs) can be found in the National Infection Prevention and Control Manual <u>http://www.nipcm.scot.nhs.uk/</u>





5. Influenza Outbreak Daily Actions Checklist

For ongoing monitoring until outbreak is under control.

Daily In	fluenza Checklist: complete daily during outbreak							
	Date:							·····
Date of first symptoms:	Completed by (initials):							
	Number of new symptomatic residents today:	_						
	Confirmed cases of influenza today:							
Total cases p								
Are any	Y/N	Y/N	Y/N	Y/N	Y/N	Y/N	Y/N	
	Number of new symptomatic staff today:	_		_	_			
Resident Placement: doors to symptomatic residents rooms a								
symptomatic and asymptomatic residents are kept apart and c	ommon areas are closed							
All residents have been told about the outbreak, advised to av								
hygiene and frequent hand hygiene, and have been given the								
Resident care: Care assessments are completed for today; ar								
Resident care: Antibiotic/antiviral prescribing for all residents								
Inter-care facility transfers (if clinically essential) have been pr								
Relatives/carers of residents discharged to home care have b								
Care Worker practices: Sufficient staff are on duty for all area								
Care Worker practices: Where possible, staff are allocated to	care for symptomatic or non symptomatic residents, but not both							
There are sufficient supplies of PPE (glove, aprons, masks), A	BHR, liquid soap, disposable towels and tissues							
There is sufficient dedicated care equipment (i.e. washbowls,								
All communal resident care equipment has been cleaned follo								
All areas are clutter free and bins have been emptied and are	easily accessible for residents to dispose of tissues							
Affected areas have been disinfected as per the enhanced cle	aning protocol (1000 ppm av cl/all frequently touched surfaces)							
Following any resident discharges, and prior to admitting a new	resident, a terminal clean of the room/area has been carried out							
Residents and visitors have been updated about any restriction	is or changes to visiting policy							
All changes to outbreak status have been communicated to s	aff/relatives and residents within the facility							
A terminal clean of the wider facility has been organised w least 48 hours	ith the HPT once all persons have been symptom-free for at							





6. Influenza Outbreak Data Record

For monitoring all residents symptoms

Complete for all symptomatic cases

(See over page for example)

Name of resident/staff	505	Had flu	Antibiotic	Antiviral	Specimen	Specimen	Date of start and end of symptoms														
member	DOB	vaccine? (Y or N)	(Y or N)	(Y or N)	date (X)	result (+ or -)															
																	-				
								1													
								1													
									l	1								1			



		Had flu vaccine? (Y or N)	Antibiotic (Y or N)	Antivirals (Y or N)		Specimen result (+ or -)		Date of start and end of symptoms															
Name of resident/ staff member	DOB				Specimen date (X)		19 /9	20 /9	21 /9		23 /9	24 /9	25 /9	26 /9	27 /9	28 /9	29 /9	30 /9	1/ 10	2/ 10	3/ 10	4/ 10	5/ 10
Joe bloggs	1/1/32	Ν	Ν	N	20/9	+	•	Х					•										
Jane Brown	5/9/44	Y	Ν	Y					•				•										



7. Influenza - Information for residents and visitors

What is influenza?

Influenza (flu) is a viral respiratory illness that can affect many people during the winter months. Symptoms include a cough (chesty or dry), runny nose, congestion, sore throat, shortness of breath, aching muscles, exhaustion, cold sweats and a fever/high temperature.

Care home residents are at increased risk of complications including secondary chest infections and pneumonia and may require admission to hospital for supportive care. In these cases, flu can be life-threatening.

What is the treatment?

For most people, plenty of rest, keeping warm, drinking plenty of fluids (to prevent dehydration) and taking over-the-counter paracetamol or ibuprofen (to lower temperature and relieve aches) will help ease the symptoms. For more vulnerable people (pregnant women, children, the elderly, those with poor health), the GP may prescribe antiviral medication.

How is it spread?

When infected people cough or sneeze, the virus is sprayed into the air and droplets land on nearby surfaces, objects and equipment. People become infected in the following ways:

- Having close contact with an infected person getting coughed or sneezed on, or shaking hands.
- Touching contaminated surfaces/objects (e.g door handles, tables, magazines).
- Eating food or drinks that someone who is ill has coughed or sneezed on.

Someone with flu can be contagious before they feel ill and can still spread the virus for up to a week after their symptoms have stopped.

What is an outbreak?

An outbreak happens when 2 or more people have flu at the same time in the same place; this can happen in hospitals, care homes and in the community.

Care home residents are extremely vulnerable because they are often elderly and frail, have underlying health conditions and are living in a closed environment where the virus can spread easily.





How do care homes control influenza?

- Staff will wear protective clothing (disposable aprons and gloves) and facemasks.
- Visitors will not be expected to wear disposable aprons and gloves unless they are helping a resident with personal care but may be offered facemasks.
- Cleaning and disinfection of affected areas will be increased.
- A restriction on visiting may be put in place if lots of residents are affected; you will be kept up-to-date of any changes to visiting policy and can ask a member of staff if you are concerned.
- Non-essential visits from hairdressers, podiatrists etc may be postponed.
- Residents may be offered preventative antiviral mediation even if they have already had their seasonal flu vaccination and even if they have not yet developed the flu.

What can visitors do to help?

- ✓ Make sure you get your seasonal **flu vaccination** (if you are eligible).
- Wash your hands frequently with soap and water; alcohol hand rubs can be used between washes.
- Make sure you practice good respiratory/cough hygiene; cover your mouth when coughing or sneezing and try and catch it with a tissue; bin the tissue immediately and wash your hands straight away.
- Wear disposable aprons, gloves and a facemask if advised by a member of staff.
- × Do not leave food open and exposed in residents rooms.
- × If you are pregnant, have small children, or have a long term health condition we advise that you do not visit during a flu outbreak for your own safety; please discuss with a member of staff.
- ✗ If you think you might have the flu, please do not visit the care home as you could spread it; you can start visiting again when you have been symptom-free for at least 48 hours.

What can residents to do help?

- Wash your hands frequently with soap and water; alcohol hand rubs can be used between washes.
- Make sure you practice good respiratory/cough hygiene; cover your mouth when coughing or sneezing and try and catch it with a tissue; bin the tissue immediately and wash your hands straight away.
- ✓ If you are feeling unwell, tell a member of staff immediately.
- If you catch the flu, staff will ask you to stay in your room and keep the door closed until you are better.
- × Do not use common areas of the home during a flu outbreak.





8. Posters for use during outbreaks





ATTENTION ALL VISITORS

We are currently experiencing an outbreak of influenza!

If you are pregnant, have a chronic health condition, or have small children, we recommend that you do not visit during this time - **please see a member of staff for more information**








Due to an outbreak of flu we are applying some infection control measures to help us stop the virus spreading.

Please see a member of staff before visiting to ensure it's safe to visit and to find out what you need to do.

If you think you have the flu, please don't visit until 48 hours after your symptoms stop – you could still pass the virus on.



All visitors please:



• Wash your hands after arriving and before leaving



 Use a tissue when you cough or sneeze



• Put used tissues in the bins provided



 Wash your hands after coughing or sneezing

If you are ill and must visit, please speak to a member of staff